

Immunosuppression in Kidney Transplant in Children & 40 kg

LD/AD	Day	Tacrolimus (PO)	MMF (PO) alt. AZA (PO)	Prednisolone (PO)	
LD	—2		600 mg/m ² x 2 3 mg/kg x 1.	10 mg/m ² x 2	
	—1		600 mg/m ² x 2 3 mg/kg x 1	10 mg/m ² x 2	
LD/AD	Op	0	0	0	0.25 g/m ² SM IV (max. 500 mg)
	+ 1	0.15 mg/kg x 2	600 mg/m ² x 2 3 mg/kg x 1	37.5 mg/m ² x 2	
	+ 2	Concentration			32.5 mg/m ² x 2
	+ 3				27.5 mg/m ² x 2
	+ 4				22.5 mg/m ² x 2
	+ 5				17.5 mg/m ² x 2
	+ 6				15 mg/m ² x 2
	+ 7				12.5 mg/m ² x 2
	+ 8				10 mg/m ² x 2
	+ 9				15 mg/m ² x1
	30				10 mg/m ²
	∨				1
	60				7.5 mg/m ²
	∨				
90			450 mg/m ² x 2 1 mg/kg x 1	Transition to every other day beh enl spec PM	

If IV give Solumedrol (dose= prednisolone x 0.8)

Tacrolimus concentration	
Month	LC-MS/MS
&1	8 ng/mL
1-3	4-8 ng/mL
> 3	4 ng/mL

CellCept AUC 95-190 μM*h



Version history

Each document should contain a history that tells you, for each version, what changed, who made the change and when the change was made.

Version	Date	Change and comment	Responsible
2	2018-12-28	New template, new organisation.	Lars Wennberg
1	2013-11-19	Document updated and transferred to Domino LIS document template	Lars Wennberg/Tarja Tervonen

